arXiv:cond-mat/0009100v1 [cond-mat.mes-hall] 7 Sep 2000

Intermediate spin, Schrödinger cat states and nano-magnets.

S. E. Barnes
Department of Physics, University of Miami, Coral Gables, Florida 33124

(February 1, 2008)

Quantum tunneling of nano-magnets finds a natural description in terms of intermediate spin. Periodic magnetic effects correspond to a change of flux by the flux quantum Φ_0. Schrödinger cat states with different superpositions of the applied magnetic field occur. The molecular magnet Fe_8 is discussed and oscillations are predicted for Mn_{12}.

75.45+j,75.50Tt,75.60Ej

Recent experiments by Wernsdoffer and Sessoli [1] on the molecular magnet known as Fe_8 show fasinating behaviour as a function of an applied field. This is an orthrombic easy axis material with one perpedicular axis harder than the other. As a function of either an easy *or* hard axis field the tunnel splitting manifests self-similar behavior at periodic values of these fields. As a function of the third component the splitting increases essentially exponentially. While these results can be well reproduced by the exact diaginalisation of a suitable large spin model, as yet there is not an analytic theory which permits a clear understanding of the observed phenomina. An adequate such theory should permit the prediction of new related effects for materials, e.g., of different symmetry.

Invariably, in the tunnelling regime, the least action paths which make any significant contribution to the tunneling amplitude can be mapped to a circle. The problem is then equivalent to the motion of a charged particle in a magetic field, Fig. (1), i.e., to the problem of *intermediate spin*. In this Letter will be described an analytic theory based upon an extension of the intermediate spin approach developed earlier by the author [2]. In terms of this the behavior of Fe_8 is easily understood and, e.g., a new periodic effect can be predicted for the large spin model often used to describe the much studied [3] Mn_{12}.

The relevant spin model for Fe_8 can written as:

$$\mathcal{H} = -(D-E){S_x}^2 + 2E{S_z}^2 \\ -ES^2 - hS_z - h_\ell S_x - h_t S_y, \qquad (1)$$

where the positive anisotropy constants have $E < D$ [4]. The magnetic field h, parallel to the hardest axis, is equivalent to that in Fig. (1) and most physical quantities, including the tunnel splitting, must be periodic in the flux Φ produced by this field, this with a period Φ_0, the flux quantum. If the physical spin value S is whole integer, a flux of $(n+\frac{1}{2})\Phi_0$ corresponds to a statistical parameter $\alpha_s = 1$, modulo unity, and transmutes the model to half-integer spin while intermediate values $n\Phi_0$ reflect a $\alpha_s = 0$, modulo unity, and are whole-integer points. Kramer's theorem implies a zero tunnel splitting for half-integer points. These facts are reflected in the quasi-periodic tunnel splitting $\Delta = \Delta_0 \cos 2\pi\Phi/\Phi_0$ with $\Phi/\Phi_0 = h/4\hbar\sqrt{2E(D+E)}$ evident in experiment [1].

There are some interesting surprises, e.g., in order to describe tunneling in Fe_8 it is necessary to construct Schrödinger cat states which involve different values of the applied magnetic field. Even more bizarre is that the other transverse field h_t is equivalent to an *imaginary* flux Φ leading to a *cosh* dependence of the splitting. In general for an arbitrarly directed transverse field the effective flux is complex!

There are classical ground states which with $h = h_t = 0$ lie along the positive and negative x-axes. Tunneling occurs between these ground states. Adding a the longitudinal field h_ℓ, e.g., the ground state for one direction can be brought to the same energy as the first excited state in the other direction. In this case one wave function has even parity and the other odd and this adds a phase shift of π to the wave function. This also converts whole into half-integer spin, again an effect seen in experiment [1].

These ideas permit the prediction of new effects even for the extensively studied molecular magnet Mn_{12}. This is a tetragonal easy y-axis magnet but with equivalent x and y hard directions in the transverse plane. There are *four* equivalent least action tunnel paths in the absence of a field. When a modest field is applied along any of the four hard directions there remain only two equivalent least action paths which are mapped to the intermediate spin circle. Actually this problem can be solved even for



FIG. 1. A particle of charge q is confined to a unit circle. A tunneling problem might be defined by adding a potential V such that $V(\phi) = V(\phi+\pi)$. If this potential has minima at $\phi = 0$ and π and the barrier between is higher than the zero point energy then there will be a ground doublet with a splitting Δ. A perpendicular field exerts no force on the particle but does change the slitting as explained in the text.



FIG. 2. The analytic result is $\Delta_0|\cosh(\pi h/p)\cos(\pi h/p)|$ where $p \approx 2S^{1/2}(12aD^3)^{1/4}$. This corresponds to a complex magnetic field in which the real and imaginary parts are equal. This exact result is indicated. The difference becomes smaller with increasing S.

small fields by mapping the four paths to *two* circles and then adding the wavefunctions in much the same way as in the elementary theory of a two slit experiment. These circles have the applied field at 45^o and the flux is complex with equal real and imaginary parts. As with Fe_8 there are periodic half-integer points at which the splitting is zero but now the maximum splitting increases by almost an order of magnitude per period. The result is given in the caption to Fig. (2) which compares it with that of exact diagonalisation. Further details will be given elsewhere [4]. It can also be shown that similar oscillations occur when the field is directed along a hard three or four fold axis of a cubic material [4].

For the basic intermediate spin problem [5] of a charged particle in two dimensions, Fig. (1), the relevant group is SO(2) which is equivalent to U(1), i.e., is abelian with a single operator S_z, the generator of rotations $e^{iS_z\phi}$, where the z-axis is perpendicular to the circle and where ϕ is the angle to the x-axis. For a unit circle $S_z = p_\phi$ the momentum conjugate to ϕ, i.e., $[\phi, p_\phi] = i\hbar$. The eigenstates of S_z and p_ϕ are $\psi_m(\phi) = \frac{1}{(2\pi)^{1/2}}e^{im\phi}$ but for particles on a circle there is no reason to insist on m being an integer reflecting the fact that the spectrum of S_z for $SO(2)$ is continuous [5]. In general for this *singular gauge* $\psi_m(\phi)$ is multi-valued. A convenient single valued function can be defined by taking some convention for $\psi_m(\phi)$ in the interval $\{0, 2\pi\}$ and then analytically continuing this function to the interval $\{-\infty, +\infty\}$. For any physical potential $V(\phi) = V(\phi + 2\pi n)$; $n = 0, \pm1, \pm2\ldots$, and implies that only the basis states $\psi_{\frac{\alpha_s}{2}+n}(\phi)$; $n = 0, \pm1, \pm2\ldots$, mix. It follows for a given statistical parameter α_s the energy spectrum is discrete. This can be stated differently: Since $V(\phi)$ is periodic the solutions obey Floquet's (Bloch's) theorem, i.e., $\psi_k(\phi) = e^{ik\phi}u_k(\phi)$ where $u_k(\phi)$ is periodic. The reciprocal lattice vector $K = 2\pi/(a = 2\pi) = 1$ and the energy $\epsilon(k) = \epsilon(k + nK)$; $n = 0, \pm1, \pm2\ldots$ i.e., is periodic. An examination of the Fourier components immediately implies $k = \alpha_s/2$. Given that $V(\phi)$ has a point with reflection symmetry the $\psi_{\pm k}(\phi)$ are degenerate and the most general solution for a given energy is $A\psi_k(\phi) + B\psi_{-k}(\phi)$ for which the spectrum of $m = n \pm \frac{\alpha_s}{2}$, i.e., there are twice the number of observable S_z values *except* importantly for the whole and half-integer points when $\alpha_s = 0$ or 1. This more general method of extrapolating between whole and half-integer spin which will be found applicable here.



FIG. 3. Mapping of the unit sphere to a two dimensional rectangle

In the *non-singular gauge* the basis set $\psi_n(\phi)$ has $n = \pm1, \pm2\ldots$ and the charged particle encircles a magnetic flux $\Phi = \alpha_s\Phi_0/2$ where Φ_0 is the appropriate flux quantum. For the same $V(\phi)$ there exists an eigenstate $u_k(\phi)$ with the same energy as in the singular gauge, i.e., the energy spectrum are identical and two problems can mapped to each other and both reflect intermediate spin. In the non-singular gauge the combination $Au_k(\phi) + Bu_{-k}(\phi)$ comprises a linear combination of states with the flux Φ in both its positive and negative senses. Evidently the expectation value of the flux $\langle\Phi\rangle$ is in general smaller than $\alpha_s\Phi_0/2$. This possibility of a linear combination of different flux states does not seem to have been envisaged in connection with intermediate statistics although it would arise if, e.g., quantum coherence in SQUIDS could be observed. It is realized in the systems, e.g., Fe_8, discussed here.

The usual functional integral formulation [6] involves coherent state $|\theta, \phi\rangle$ obtained by rotating $|S_z = S\rangle$ through usual angles θ, ϕ. *However* the $|\theta, \phi\rangle$ are *very* much over-complete. In fact the states $|\phi\rangle \equiv |\theta = \frac{\pi}{2}, \phi\rangle = (2S+1)^{1/2}\sum_{m=-S}^{S} e^{-im\phi}|m\rangle$, where $|m\rangle \equiv |S_z = m\rangle$, remain over-complete. If $|\psi\rangle = \sum_{m=-S}^{S} a(m)|m\rangle$ then $\psi(\phi) \equiv \langle\phi|\psi\rangle = \sum_{m=-S}^{S} e^{im\phi}a(m)$, i.e., is a Fourier transform of the m-space wave function $a(m)$. The function $\psi(\phi)$ is defined on the interval $\{-\infty, +\infty\}$ and the inverse transform $\psi(m) \equiv \frac{1}{2\pi}\int_{-\infty}^{+\infty} d\phi\psi(\phi) = \sum_{m'}\delta(m - m')a(m')$ so that $|\psi\rangle = \int_{-\infty}^{+\infty} dm\,\psi(m)|m\rangle$, i.e., $\psi(m)$ is a definition of the m-space wave function appropriate when m is considered as a continuous variable. Clearly $\hat{S}_z\psi(\phi) = \int_{-\infty}^{+\infty} dme^{im\phi}\hbar m\psi(m) = p_\phi\psi(\phi)$ where $p_\phi = -i\hbar\frac{d}{d\phi}$ is the same as that defined earlier. Also evident is that multiplication by $e^{i\phi}$ amounts to a displacement of m by unity and so, e.g., $\hat{S}^+ = e^{i\phi}[\hbar^2 S(S+1) - p_\phi(p_\phi + 1)]^{1/2}$. These definitions permit an arbitrary spin Hamiltonianto be expressed in terms of the cononical p_ϕ and ϕ.

In terms of the unit sphere ϕ and p_ϕ might be considered as a particular projection in which the axis of quantization plays the role of the North-South axis as shown in Fig. (3). In these terms, the angle ϕ is evidently the longitude while p_ϕ is the projection on the z-axis and is therefore the sine of the latitude.

It is necessary to carefully formulate the problem in terms of the basis set $|m\rangle$, i.e., in m-space. If $|\psi\rangle = \sum_{m=-S}^{+S} a(m)|m\rangle$, then in this basis Schrödinger's equation for Eqn. (1) can be written as ($h_\ell = h_t = 0$):

$$\begin{aligned}&\left(\epsilon - (2E\hbar^2 m^2 - h\hbar m)\right) a(m)\\ &= \tfrac{1}{4}\hbar^2(D-E)\Big[M_m^{m+1}M_{m+1}^{m+2}a(m+2)\\ &\quad + M_m^{m-1}M_{m-1}^{m-2}a(m-2)\\ &\quad + [M_m^{m+1\,2} + M_m^{m-1\,2}]a(m)\Big],\end{aligned} \tag{2}$$

where $M_s^t = [S(S+1) - st]^{1/2}$. The structure of this equation is worth noting. It has the form of *two* tight binding models, one for even m values and the other for odd m.

The wave function for the eigenstate $|S_x = S\rangle$ is $a^0(m) = \frac{1}{2^S}\left[\frac{(2S)!}{(S-m)!(S+m)!}\right]^{1/2}$ and $a^0(m)$ has to be a solution to Eqn. (2) with $h = E = 0$, this corresponding to the absence of tunneling. Using Stirling's approximation this reduces to $a^0(m) \propto e^{-m^2/2S}$, however the deviations from this approximation are very important. The absence of tunneling when $h = E = 0$ is put in evidence by a transformation: $a(m) = a^0(m)f(m)$. With the definition $f(\phi) = \sum_m e^{im\phi}f(m)$ this yields the *exact* Schrödinger's equation:

$$\Big(\mathcal{E} - 2E{p_\phi}^2\Big)f(\phi) = V(\phi, p_\phi)f(\phi) \tag{3}$$

with

$$\begin{aligned}V(\phi, p_\phi) &= \tfrac{1}{2}(D-E)\times\\ \Big[2\sin^2\phi(\hbar^2 S(S-1) &+ {p_\phi}^2) - i\hbar(2S-1)\sin 2\phi p_\phi\Big],\end{aligned} \tag{4}$$

where $\mathcal{E} = (\epsilon - E^0)$; $E^0 = -\hbar^2 S^2(D+E))$. This is of the intermediate spin form but with a momentum dependent potential, $V(\phi, p_\phi)$.

A new wave function is defined by $f(\phi) = (1 - (\alpha_s p_\phi/2\hbar S))g(\phi)$. This admixes, e.g., a little of the first excited into the ground state. If $g(\phi)$ is the solution of

$$\begin{aligned}\Big(\mathcal{E} - 2E\left(p_\phi + \hbar\tfrac{\alpha_s}{2}\right)^2 + \hbar^2 E\frac{{\alpha_s}^2}{2}\Big)g(\phi)&\\ = V(\phi, p_\phi + \hbar\frac{\alpha_s}{2})g(\phi),&\end{aligned} \tag{5}$$

then for modestly large S, $f(\phi) = (1 - (\alpha_s p_\phi/2\hbar S))g(\phi)$ is the solution to

$$\begin{aligned}\Big(\mathcal{E} - 2E{p_\phi}^2 + 2\hbar(D+E)\frac{\alpha_s}{2}p_\phi\Big)f(\phi)&\\ = V(\phi, p_\phi)f(\phi).&\end{aligned} \tag{6}$$

The shift in p_ϕ determines the wave vector: $k = \alpha_s/2$. In turn the statistical parameter must be chosen to give ground state energy determined without tunneling. Required is that $\hbar^2 E\frac{{\alpha_s}^2}{2} = \frac{h^2}{8(D+E)}$, so $k = \frac{\alpha_s}{2} = h/2\hbar\sqrt{2E(D+E)}$. This fixes the Zeeman term in Eqn. (6); $2\hbar(D+E)\frac{\alpha_s}{2}p_\phi = \sqrt{(D+E)/2E}hp_\phi$. Since, in general, this is larger than hp_ϕ the solution is a Schrödinger cat state, $A(1 - (\alpha_s p_\phi/2\hbar S))u_k(\phi) + B(1 + (\alpha_s p_\phi/2\hbar S))u_{-k}(\phi)$ involving a linear combination of both bare fields $\pm\sqrt{(D+E)/2E}h$ such that the expectation value is equal the physical applied field. This requires $A = 1 + \sqrt{2E/(D+E)}$ and $B = 1 - \sqrt{2E/(D+E)}$. The total energy for level n and wave vector k is,

$$E_n(k) = E^0 - \frac{h^2}{8(D+E)} + \left(n + \frac{1}{2}\right)\hbar\omega_0 + \epsilon_n(k). \tag{7}$$

The first two terms are purely classical in origin while the third corresponds to the harmonic motion about the bottom of the well. It is the last term which reflects the tunneling between wells.

The problem therefore reduces to the determination of $\epsilon_n(k)$ and when the tunneling approximation is well justified this reflects the tight binding approximation. With only a transverse field h, the problem has a higher symmetry and potential $V(\phi, p_\phi)$ has a ϕ period of $a = \pi$ rather than 2π so the reciprocal space unit vector $K = 2$. For a tight binding model the tunneling energy, $\epsilon_n(k) = \frac{\Delta_0}{2}\cos\pi k$ where $\Delta_0/2$ is the undetermined matrix element for tunneling for level n. (The method for determining this is given in [4].) There are *two* solutions which are acceptable for $h = 0$, namely $k = \frac{\alpha_s}{2} = 0$ corresponding to even m values *and* $k = \frac{\alpha_s}{2} = 1$ and odd m. These two k values correspond to $\epsilon(k) = \pm\frac{\Delta_0}{2}$ and so the zero field tunnel splitting is reflected in the parameter Δ_0. For finite transverse fields h the corresponding $k = \frac{\alpha_s}{2} = (h/2\hbar\sqrt{2E(D+E)})$ and $k = 1 + \frac{\alpha_s}{2} = 1 + (h/2\hbar\sqrt{2E(D+E)})$ and so the tunnel splitting is,

$$\Delta = \Delta_0\cos\frac{\pi h}{2\hbar\sqrt{2E(D+E)}}, \tag{8}$$

i.e., of the form mentioned in the introductory remarks. An equivalent result was first obtained some time ago by Garg [7] and attributed to "topological quenching".

Nowhere in the development has use been made of the assumption that S is a whole-integer. The only difference when S is a half-integer is that the values of m are also half-integer and thus the solutions correspond to $k = \frac{1}{2} + \frac{\alpha_s}{2} = \frac{1}{2} + (h/2\hbar\sqrt{2E(D+E)})$ and $k = -\frac{1}{2} + \frac{\alpha_s}{2} = -\frac{1}{2} + (h/2\hbar\sqrt{2E(D+E)})$. This has the effect of replacing the cosine is replaced by a sine.

Consider first the effect of adding h_ℓ alone. This adds a potential, i.e., $V(\phi, p_\phi) \Rightarrow V(\phi, p_\phi) + \hbar S h_\ell \cos\phi$ and removes the symmetry between $\phi = 0$ and π. The period of the potential is now the full $a = 2\pi$, with $K = 1$, *but* with two wells per unit cell. The harmonic levels near $\phi = \pi$ have quantum numbers designated by n and are higher than those near $\phi = 0$ with labels n'. Ignoring tunneling these have energies, $E_n = E_0 - \frac{h^2}{8(D+E)} + \hbar n h_\ell + \left(n + \frac{1}{2}\right)\hbar\omega_0$, and, $E_{n'} = E_0 - \frac{h^2}{8(D+E)} - \hbar n' h_\ell + \left(n' + \frac{1}{2}\right)\hbar\omega_0$. For a given h there is only one solution with $k = \frac{\alpha_s}{2} = (h/2\hbar\sqrt{2E(D+E)})$. There are now two possibilities (i) $E_n \neq E_{n'}$ and there is a *very* narrow band formed about both these energies. The energy differences have a *very* small dependence on k and hence h, see below. (ii) When $E_n \approx E_{n'}$ there is resonant tunneling. There are again two cases (iia) when both n and n' are odd or even and (iib) when one of n and n' is odd and the other even. As illustrated in Fig. (4) there are two tunneling matrix elements one $\Delta_{n,n'}$ for the barrier between $\phi = 0$ and π and $\Delta_{n',n}$ for that between $\phi = -\pi$ and 0 (or π and 2π). With first $h = (h_t =)0$, Schrödinger's equation admits real solutions, $\psi(m)$ or $\psi(\phi)$, and so these matrix elements can be made real. For case (iia) when the symmetry of the wave functions is the same the tunneling matrix elements $\Delta_{n,n'} = \Delta_{n',n} \equiv \Delta_0$. For finite h, for the Floquet (Bloch) wave function, the phase advances by $2\pi k$ from one cell to the next while to agree with the smaller unit cell appropriate when $h_\ell = 0$ the phase should advance by πk from one well to the next within the cell. This leads to $\Delta_{n,n'} = e^{i\pi k}\Delta_0$ and $\Delta_{n',n} = e^{-i\pi k}\Delta_0$.

At this point it is possible to introduce the second transverse field h_t by slight of hand. It is observed that if E changes sign then the two transverse axes change role and thus h becomes the equivalent of h_t. If $\mathcal{H}$ is considered to be a function of a complex parameter E the solution might be analytically continued from positive to negative values. In particular $k = (h/2\hbar\sqrt{2E(D+E)})$ becomes pure imaginary, i.e., $k \to ik_t = i(h_t/2\hbar\sqrt{2E(D-E)})$ and it follows that $\Delta_{n,n'} = e^{i\pi(k+ik_t)}\Delta_0$ with $\Delta_{n',n} = e^{-i\pi(k+ik_t)}\Delta_0$. This information can be represented by a two level model:

$$\mathcal{H} = (E_n - E_{n'})S_z + \frac{\Delta_0}{2}\left[e^{i\pi(k+ik_t)}S^+ + e^{-i\pi(k+ik_t)}S^-\right],$$

where the center of gravity energy $\frac{1}{2}(E_n + E_{n'})$ has been dropped. Degenerate perturbation theory then yields, $\epsilon_k = \pm\frac{1}{2}\sqrt{(E_n - E_{n'})^2 + 4{\Delta_0}^2|\cos\pi(k+ik_t)|^2}$ or at resonance $\epsilon_k = \pm\frac{\Delta_0}{2}|\cos\pi(k+ik_t)|$, with $k = \frac{h}{2\hbar\sqrt{2E(D+E)}}$ and $k_t = \frac{h_t}{2\hbar\sqrt{2E(D-E)}}$. The $h_\ell = h_t = 0$ result is recovered when $n = n'$. When for (iib) one wave function changes sign while the other does not it is necessarily the case, for $h = h_t = 0$, that $\Delta_{,n'} = -\Delta_{n',n} \equiv \Delta_0$ whence everywhere in ϵ_k the cosine becomes a sine and the result is as if the original S was half-integer! This explains the alternation observed in experiment [1].



FIG. 4. When, e.g., $n = 0$ and $n' = 2$, both wave functions are even there is no sign change. If $n = 0$ and $n' = 1$ the wave must change sign as its ϕ argument advances by 2π.

Certainly the most striking claim is that the wave function $A(1-(\alpha_s p_\phi/2\hbar S))u_k(\phi) + B(1+(\alpha_s p_\phi/2\hbar S))u_{-k}(\phi)$ for an intermediate spin system is a mixture of a state with the local applied field of magnitude $h_p = \pm\sqrt{(D+E)/2E}h$, for small fields. The fields are both parallel and anti-parallel to the distant applied field h, but see [4]. (With an equivalent claim for h_t which involves $\sqrt{(D-E)/2E}h_t$.) *Experimentally* the existence of this superposition is easy to establish. The transverse field h couples to the physical S_z, i.e., this operator in the non-singular gauge, and which has a spectrum with a period $\hbar$ or in units of magnetic moment $g\mu_B\hbar$ where g is the g-factor. Thus if M_z is the measured moment in the direction of the transverse field, $m = M_z/g\mu_B\hbar$, the dimensionless magnetization has a period of unity. This would be the case if the state did not consist *precisely* of a superposition of field directions. Because of the admixture, the measurable period is reduced to $\Delta m \equiv m_1 = \sqrt{\frac{2E}{D+E}} \leq 1$ which is confirmed by exact results but remains to be tested experimentally.

---

[1] W. Wernsdorfer and R. Sessoli, Science **284**, 133 (1999).
[2] S.E. Barnes, J. Phys. Cond. Matter 10, L665 (1998); and cond-mat/9907257.
[3] L. Thomas, F. Lionti, R. Ballou, R. Sessoli, A. Caneschi and B. Barbara, Nature **383**, 145 (1996); J. R. Friedman et al., Phys. Rev. Lett. **76**, 3820 (1996).
[4] S.E. Barnes, unpublished, cond-mat/00XXXX.
[5] See e.g., A. Khare *Fractional statistics and quantum field theory*, World Scientific, Singapore (1997).
[6] E. M. Chudnovsky and J. Tejada, *Macroscopic Quantum Tunneling of the Magnetic Moment* (Cambridge University Press, 1997).
[7] A. Garg, Europhys. Lett. **22**, 205 (1993).